



GAPDH (A531) polyclonal antibody

Catalog:: QS04019

BackGround:

Glyceraldehyde 3 phosphate dehydrogenase (GAPDH) is well known as one of the key enzymes involved in glycolysis. As well as functioning as a glycolytic enzyme in cytoplasm, recent evidence suggests that mammalian GAPDH is also involved in a great number of intracellular proceses such as membrane fusion, microtubule bundling, phosphotransferase activity, nuclear RNA export, DNA replication, and DNA repair. During the last decade a lot of data appeared concerning the role of GAPDH in different pathologies including prostate cancer progression, programmed neuronal cell death, age related neuronal diseases, such as Alzheimer's and Huntington's disease. GAPDH is expressed in all cells. It is constitutively expressed in almost all tissues at high levels. There are however some physiological factors such as hypoxia and diabetes that increase GAPDH expression in certain cell types. GAPDH molecule is composed of four 36kDa subunits.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide,50% glycerol, pH7.2

Molecular Weight:

~ 36 kDa

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE)..

Applications:

WB:1:5000~1:20000


Storage&Stability:

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Specificity:

GAPDH (1A6) mAb detects endogenous levels of GAPDH protein.

DATA:



地址: 江苏省镇江市宝华镇仙林东路16号双创大厦1207室

www.qianzhusong.com 1 / 2




江苏千株松生物科技有限公司



Western blot (WB) analysis of GAPDH (A531) polyclonal antibody at 1:5000-1:20000 dilution Lane1:The Kidney tissue lysate of Mouse(40ug) Lane2:The Kidney tissue lysate of Rat(40ug) Lane3:PC12 whole cell lysate(40ug) Lane4:CT26 whole cell lysate(40ug) Lane5:A549 whole cell lysate(40ug)

Lane6:L02 whole cell lysate(40ug)



Immunofluorescence analysis of RAW264.7 cells using GAPDH pAb at dilution of 1:1000 (40x lens). **Note:**

For research use only, not for use in diagnostic procedure.

2 / 2