



JNK1/2/3 (phospho-T183) polyclonal antibody

Catalog: QS04123

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

The human JNK1 (PRKM8, SAPK1, MAPK8) gene maps to chromosome 10q11.22 and shares 83% amino acid identity with JNK2. JNK1 is necessary for normal activation and differentiation of CD4 helper T (TH) cells into TH1 and TH2 effector cells. Capsaicin activates JNK1 and p38 in Ras-transformed human breast epithelial cells. Nitrogen oxides (NO_x) upregulate JNK1 in addition to c-Fos, c-Jun and other signaling kinases, including MEKK1 and p38. JNK3 (MK10, MAPK10, PRKM10) is activated by pro-inflammatory cytokines and environmental stress by phosphorylating transcription factors such as c-Jun and ATF2. This is important for AP-1 transcriptional activity regulation. JNK3 is crucial for neuronal apoptosis (stress-induced).

Product:

Rabbit IgG in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 54 kDa

Swiss-Prot:

P45983/P45984/P53779

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:250~1:500

IHC: 1:25~1:100

IF: 1:25~1:100


Storage&Stability:

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Specificity:

p-JNK1/2/3 (T183) polyclonal antibody detects endogenous levels of JNK1/2/3 protein only when phosphorylated at Thr183.

DATA:



Western blot (WB) analysis of p-JNK1/2/3 (T183) pAb at 1:500 dilution


Lane1:HEK293T whole cell lysate(40ug)

Lane2:MCF-7 whole cell lysate(40ug)

Lane3:K562 whole cell lysate(20ug)

Lane4:The Brain tissue lysate of Rat(40ug)

Lane5:The Kidney tissue lysate of Mouse(20ug)



Immunohistochemistry (IHC) analyzes of p-JNK1/2/3 (T183) pAb in paraffin-embedded human colorectal carcinoma tissue at 1:50.

Note:

For research use only, not for use in diagnostic procedure.